

The benefits and risks of non-steroidal anti-inflammatory drugs

Poonam Ruhel, Palika Sehgal

Department of Pharmacy, Panipat Institute of Engineering and Technology, Panipat, Haryana, India

Correspondence:

Poonam Ruhel, Department of Pharmacy, Panipat Institute of Engineering and Technology, Samalkha, Panipat - 132 102, Haryana, India.
E-mail: ruhalpoonam@gmail.com

How to cite this article: Ruhel P, Sehgal P. The benefits and risks of non-steroidal anti-inflammatory drugs. Innov Pharm Planet 2023;11(4):62-64.

Source of Support: Nil.

Conflicts of Interest: None declared.

Date of Submission: 18-10-2023

Date of Revision: 02-11-2023

Date of Acceptance: 12-11-2023

ABSTRACT

Non-steroidal anti-inflammatory drugs (NSAIDs) are commonly utilized for their analgesic, anti-inflammatory, and antipyretic properties, making them a cornerstone in managing pain and inflammation across various clinical settings. This review provides a comprehensive examination of the benefits and risks associated with NSAID use. NSAIDs function by inhibiting cyclooxygenase enzymes, which play a pivotal role in prostaglandin synthesis, thereby alleviating pain and reducing inflammation. Their effectiveness in treating conditions such as osteoarthritis, rheumatoid arthritis, and post-operative pain is well-documented. However, the benefits come with notable risks, including gastrointestinal complications such as dyspepsia, ulcers, and bleeding, as well as cardiovascular issues, such as heart attack and stroke. In addition, renal and hepatic adverse effects and allergic reactions pose significant concerns. This review aims to balance the therapeutic advantages of NSAIDs against their potential risks, emphasizing the importance of individualized treatment plans.

Keywords: Cardiovascular effects, gastrointestinal risks, non-steroidal anti-inflammatory drugs, pain relief, safety guidelines

Introduction

Non-steroidal anti-inflammatory drugs (NSAIDs) are widely used medications that help alleviate pain, reduce inflammation, and lower fever. They are commonly prescribed for various conditions, including arthritis, muscle pain, and other inflammatory disorders. NSAIDs work by inhibiting the cyclooxygenase (COX) enzymes, which play a crucial role in the synthesis of prostaglandin compounds that mediate inflammation and pain. While NSAIDs are effective in managing symptoms, they also come with a range of potential side effects and risks that must be carefully considered.^[1]

The purpose of this review is to assess the benefits and risks associated with NSAIDs. On one hand, these medications provide significant relief from pain and inflammation, making them a first-line treatment option for many patients. Clinical studies have shown that NSAIDs, such as

ibuprofen and naproxen are more effective than acetaminophen in managing pain associated with conditions such as osteoarthritis. However, the use of NSAIDs is not without complications. They are associated with gastrointestinal (GI) side effects, including dyspepsia, ulcers, and bleeding, particularly with long-term use or in patients with pre-existing GI conditions. Moreover, recent findings have highlighted an increased risk of cardiovascular events, such as heart attacks and strokes, associated with certain NSAIDs, particularly those that are COX-2 selective.

Given these considerations, it is essential for healthcare providers to weigh the benefits of NSAID therapy against the potential risks, tailoring treatment plans to individual patient needs and risk factors. This review aims to provide a comprehensive overview of the therapeutic advantages of NSAIDs while also addressing the safety concerns that accompany their use, ultimately guiding informed decision-making in clinical practice.^[2]

Mechanism of Action

The mechanism of action of NSAIDs involves both pharmacodynamics and pharmacokinetics, which are critical to understanding how these medications alleviate pain and inflammation.

Access this article online

Website: <https://innovationaljournals.com/index.php/ip> **e-ISSN:** 2348-7275

DOI: 10.31690/ipplanet.2023.v01i04.017

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution NonCommercial Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Pharmacodynamics

NSAIDs primarily exert their effects through the inhibition of COX enzymes, which are crucial in the biosynthesis of prostaglandins. There are two main isoforms of COX: COX-1 and COX-2. COX-1 is constitutively expressed in many tissues and is involved in maintaining normal physiological functions, such as protecting the gastric mucosa and regulating renal blood flow. In contrast, COX-2 is inducible and is primarily expressed at sites of inflammation.

By inhibiting these enzymes, NSAIDs decrease the production of prostaglandins, which are responsible for promoting inflammation, pain, and fever. This reduction in prostaglandin synthesis leads to the analgesic (pain-relieving), anti-inflammatory, and antipyretic (fever-reducing) effects of NSAIDs. However, the inhibition of COX-1 can lead to GI side effects, such as ulcers and bleeding, highlighting the importance of the balance between therapeutic effects and adverse reactions.

Pharmacokinetics

The pharmacokinetics of NSAIDs encompass their absorption, distribution, metabolism, and excretion. NSAIDs are generally well absorbed from the GI tract, with most achieving rapid and complete absorption after oral administration. However, the bioavailability of some NSAIDs, such as aspirin, can be lower due to hydrolysis in the GI tract or during first-pass metabolism in the liver.^[3]

Once absorbed, NSAIDs are highly bound to plasma proteins, particularly albumin, with binding rates often exceeding 90%. This high protein binding limits their distribution to body tissues, resulting in a low volume of distribution, typically ranging from 0.1 to 0.3 L/kg. The pharmacokinetics can be influenced by various factors, including age, disease state, and interactions with other medications.

NSAIDs are primarily metabolized in the liver through oxidation and conjugation to form inactive metabolites, which are then excreted in urine or bile. The elimination half-lives of NSAIDs can vary widely, from a few hours to over 70 h, depending on the specific drug and its formulation. For example, ibuprofen has a half-life of approximately 2–4 h, while naproxen can range from 12 to 17 h. Renal excretion plays a minor role in the elimination of most NSAIDs, with the exception of a few, such as azapropazone, which may be excreted unchanged.^[4]

Benefits of NSAIDs

NSAIDs are widely used medications known for their effectiveness in treating pain and inflammation. They are commonly prescribed for various conditions, including osteoarthritis, rheumatoid arthritis, and post-operative pain. Understanding the benefits of NSAIDs is essential for optimizing their use in clinical practice.

Pain relief

NSAIDs are highly effective in providing relief from various types of pain. They are commonly used to manage pain associated with

osteoarthritis and rheumatoid arthritis, where they help reduce joint pain and stiffness. Clinical studies have demonstrated that NSAIDs can significantly improve pain levels and function in patients suffering from these chronic conditions. In addition, NSAIDs are often utilized for acute pain management, including post-operative pain and pain from injuries. For instance, medications such as ibuprofen and naproxen are frequently recommended for their analgesic properties, allowing patients to achieve better pain control and improved quality of life.^[5]

Anti-inflammatory effects

Beyond pain relief, NSAIDs play a crucial role in reducing inflammation and swelling in chronic conditions. By inhibiting the COX enzymes responsible for prostaglandin synthesis, NSAIDs effectively decrease the inflammatory response in the body. This anti-inflammatory effect is particularly beneficial for patients with inflammatory disorders, as it can lead to reduced swelling, improved mobility, and overall better management of symptoms. The ability of NSAIDs to target inflammation makes them a valuable component of treatment regimens for conditions, such as arthritis and other musculoskeletal disorders.

Additional benefits

In addition to their primary roles in pain relief and inflammation reduction, NSAIDs may offer potential benefits beyond these effects. For example, low-dose aspirin, a type of NSAID, has been shown to provide cardiovascular protection by inhibiting platelet aggregation, thereby reducing the risk of heart attacks and strokes in high-risk patients. This protective effect is attributed to the drug's ability to block the COX-1 enzyme, which plays a role in platelet function.

Moreover, ongoing research is exploring the potential neuroprotective effects of certain NSAIDs, suggesting that they may have roles in managing neuroinflammatory conditions. While more studies are needed to fully understand these additional benefits, the existing evidence highlights the versatility of NSAIDs in treating a range of health issues.^[6]

Risks and Side Effects

NSAIDs are widely used for their analgesic and anti-inflammatory properties; however, their use is associated with several risks and side effects that can significantly impact patient health.

GI complications

One of the most significant risks associated with NSAID use is GI complications. These can include dyspepsia, gastritis, ulcers, and severe bleeding. NSAIDs inhibit the COX enzymes, particularly COX-1, which plays a protective role in maintaining the gastric mucosal barrier. The reduction in protective prostaglandins due to COX inhibition can lead to mucosal injury and increase the risk of gastric and duodenal ulcers. Studies indicate that up to 30% of regular NSAID users may develop endoscopic lesions, including ulcers and erosions, with serious complications occurring in 1–2% of users during treatment. These complications can manifest as overt

bleeding, which may present as melena or hematemesis, and can be life-threatening, especially in older adults or those with pre-existing conditions. The risk of developing these complications is heightened in patients taking high doses of NSAIDs, those with a history of ulcers, and individuals over the age of 60.

Cardiovascular risks^[7]

NSAIDs are also associated with cardiovascular risks, including an increased likelihood of heart attack and stroke. The risk varies among different NSAIDs, with non-selective NSAIDs such as ibuprofen and diclofenac linked to a higher incidence of cardiovascular events. In contrast, naproxen is often considered to have a more neutral cardiovascular risk profile. The mechanism behind these risks is thought to involve the inhibition of COX-2, which plays a role in vascular function and platelet aggregation. Consequently, the use of NSAIDs, particularly in patients with existing cardiovascular conditions or risk factors, necessitates careful consideration and monitoring.

Renal and hepatic issues

NSAIDs can adversely affect kidney function, leading to conditions such as acute kidney injury, especially in patients with pre-existing renal impairment or those who are dehydrated. The inhibition of prostaglandins can reduce renal blood flow and glomerular filtration rate, resulting in potential renal complications. In addition, while less common, NSAIDs can also impact liver function, particularly in patients with existing liver disease or those taking other hepatotoxic medications. Regular monitoring of kidney and liver function is recommended for patients on long-term NSAID therapy to mitigate these risks.

Allergic reactions

Hypersensitivity reactions are another concern with NSAID use. Patients may experience allergic responses, ranging from mild skin

rashes to severe anaphylactic reactions. These reactions can occur due to the body's immune response to the drug or its metabolites. Patients with a history of asthma or allergic conditions may be at a higher risk of experiencing NSAID-induced hypersensitivity. It is essential for healthcare providers to evaluate patients' allergy histories before prescribing NSAIDs and to monitor for any signs of allergic reactions during treatment.^[8]

Conclusion

NSAIDs offer significant relief for pain and inflammation but come with risks, such as GI, cardiovascular, renal, and hepatic complications. A balanced approach, incorporating personalized treatment and careful monitoring, is essential to maximize benefits and minimize risks, ensuring patient safety and effective outcomes.

References

1. Fendrick AM, Greenberg BP. A review of the benefits and risks of nonsteroidal anti-inflammatory drugs in the management of mild-to-moderate osteoarthritis. *Osteopath Med Prim Care* 2009;3:1.
2. Day RO, Graham GG. Non-steroidal anti-inflammatory drugs (NSAIDs). *BMJ* 2013;346:f3195.
3. Yaksh TL, Dirig DM, Malmberg AB. Mechanism of action of nonsteroidal anti-inflammatory drugs. *Cancer Invest* 1998;16:509-27.
4. Vane JR, Botting RM. Mechanism of action of nonsteroidal anti-inflammatory drugs. *Am J Med* 1998;104:2S-8.
5. Peura DA, Goldkind L. Balancing the gastrointestinal benefits and risks of nonselective NSAIDs. *Arthritis Res Ther* 2005;7:S7-13.
6. Pisano M. NSAIDs: Balancing the risks and benefits. *US Pharm* 2016;41:24-6.
7. Wallace JL, Chin BC. New generation NSAIDs: The benefits without the risks? *Drugs Today* 1997;33:371-8.
8. Whelton A, Hamilton CW. Nonsteroidal anti-inflammatory drugs: Effects on kidney function. *J Clin Pharmacol* 1991;31:588-98.